arXiv:math/0605087v1 [math.AG] 3 May 2006

On Poincaré series of filtrations on equivariant functions of two variables [*]

A. Campillo F. Delgado S.M. Gusein-Zade [†]

Abstract

Let a finite group G act on the complex plane $(\mathbb{C}^2, 0)$. We consider multi-index filtrations on the spaces of germs of holomorphic functions of two variables equivariant with respect to 1-dimensional representations of the group G defined by components of a modification of the complex plane $\mathbb{C}^2$ at the origin or by branches of a G-invariant plane curve singularity $(C, 0) \subset (\mathbb{C}^2, 0)$. We give formulae for the Poincaré series of these filtrations. In particular, this gives a new method to obtain the Poincaré series of analogous filtrations on the rings of germs of functions on quotient surface singularities.

Introduction

In [1], [6] (see also [2] for another proof of the result of [1]) there were computed the Poincaré series (in several variables) of the so called divisorial (multi-index) filtrations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ of germs of holomorphic functions on the complex plane $\mathbb{C}^2$ at the origin and of the filtration defined by orders of a function on branches of a germ of a plane curve $(C, 0) \subset (\mathbb{C}^2, 0)$. In particular it was shown that the Poincaré series of the latter filtration coincides with the (multi-variable) Alexander polynomial of the (algebraic) link $(S^3_\varepsilon, C \cap S^3_\varepsilon)$ corresponding to the curve $(C, 0)$. Here we give versions of these results for the equivariant case, i.e. when there is an action (a representation) of a finite group G on the complex plane $\mathbb{C}^2$ and the corresponding filtrations are considered on subspaces of germs of functions equivariant with respect to 1-dimensional representations $G \to \mathbb{C}^* = \mathbf{GL}(\mathbb{C}, 1)$ of the group G. There is a problem to

---

[*] Math. Subject Class. 14B05, 16W70, 16W22.

[†] Partially supported by the grant MCYT: MTM2004-00958. The third author partially supported by the grants RFBR–04–01–00762, NSh–4719.2006.1.

define an equivariant version of the monodromy zeta function. One reason to study "equivariant" Poincare series of filtrations is the hope that they may give a hint for such a definition (at least for curve singularities).

For a finite group G, the set G^* of 1-dimentional representations $G \to \mathbb{C}^* = \mathbf{GL}(\mathbb{C}, 1)$ of the group G is a group itself. Let $R(G)$ be the ring of (complex virtual) representations of the group G and let $R_1(G)$ be its subring generated by 1-dimensional representations. Elements of the ring $R_1(G)$ are formal sums $\sum n_\alpha \alpha$ over all different 1-dimensional representations α of the group G, $n_\alpha \in \mathbb{Z}$. The multiplication is defined by the tensor product. The rings $R(G)$ and $R_1(G)$ coincide iff the group G is abelian.

Let $(V, 0)$ be a germ of a complex analytic variety with an action of a finite group G. The group G acts on the ring $\mathcal{O}_{V,0}$ of germs of functions on $(V, 0)$ by the formula $g^*f(x) = f(g^{-1}(x))$ ($f \in \mathcal{O}_{V,0}$, $g \in G$, $x \in (V, 0)$). Let $\alpha : G \to \mathbb{C}^*$ be a 1-dimensional representation of the group G

Definition: A germ $f \in \mathcal{O}_{V,0}$ is *equivariant* with respect to the representation α if $g^*f = \alpha(g) \cdot f$ for $g \in G$.

The set of germs of functions on $(V, 0)$ equivariant with respect to the representation α is a vector subspace $\mathcal{O}^\alpha_{V,0} \subset \mathcal{O}_{V,0}$. If $\alpha_1 \neq \alpha_2$, one has $\mathcal{O}^{\alpha_1}_{V,0} \cap \mathcal{O}^{\alpha_2}_{V,0} = \{0\}$. Product of functions equivariant with respect to representations α_1 and α_2 is a function equivariant with respect to the representation $\alpha_1 \cdot \alpha_2$. If the group G is abelian, $\mathcal{O}_{V,0} = \bigoplus\limits_\alpha \mathcal{O}^\alpha_{V,0}$.

For a vector space A (finite or infinite dimensional, e.g., for $\mathcal{O}_{V,0}$ or $\mathcal{O}^\alpha_{V,0}$), a 1-index (decreasing) filtration on A is defined by a function $v : A \to \mathbb{Z}_{\geq 0} \cup \{\infty\}$, such that $v(f_1 + f_2) \geq \min\{v(f_1), v(f_2)\}$, $f_i \in A$, $i = 1, 2$. An s-index filtration on A is defined by s such functions v_i, $i = 1, \ldots, s$. For $\underline{v} = (v_1, \ldots, v_s) \in \mathbb{Z}^s$, the corresponding subspace $J(\underline{v})$ is defined as $\{a \in A : v_i(a) \geq v_i, i = 1, \ldots, s\}$. (To describe the filtration it is sufficient to define the subspaces $J(\underline{v})$ only for $\underline{v} \in \mathbb{Z}^s_{\geq 0}$ (i.e., $v_i \geq 0$ for $i = 1, \ldots, s$), however it is convenient to suppose $J(\underline{v})$ to be defined for all $\underline{v}$ from the lattice $\mathbb{Z}^s$.)

Suppose that all the factor spaces $J(\underline{v})/J(\underline{v} + \underline{1})$ ($\underline{1} = (1, \ldots, 1)$, $\underline{v} \in \mathbb{Z}^s$) are finite dimensional. (This is equivalent to say that $\dim A/J(\underline{v})$ is finite for any $\underline{v} \in \mathbb{Z}^s_{\geq 0}$.) Let $d(\underline{v}) := \dim J(\underline{v})/J(\underline{v} + \underline{1})$, $\underline{v} \in \mathbb{Z}^r$. The *Poincaré series* of the multi-index filtration $\{J(\underline{v})\}$ is the series $P(t_1, \ldots, t_s) \in \mathbb{Z}[[t_1, \ldots, t_s]]$ defined by the formula

$$P(\underline{t}) = \frac{\left(\sum\limits_{\underline{v} \in \mathbb{Z}^s} d(\underline{v}) \cdot \underline{t}^{\underline{v}}\right) \prod\limits_{i=1}^{s} (t_i - 1)}{\underline{t}^{\underline{1}} - 1}\,, \tag{1}$$

where $\underline{t}^{\underline{v}} = t_1^{v_1} \cdot \ldots \cdot t_s^{v_s}$ ($\underline{t} = (t_1, \ldots, t_s)$, $\underline{v} = (v_1, \ldots, v_s)$); see, e.g., [5], [1].

Remark. Pay attention that the sum in the numerator is over all $\underline{v} \in \mathbb{Z}^s$, not only over those from $\mathbb{Z}^s_{\geq 0}$. This sum contains monomials with negative powers of variables (for $s \geq 2$), however, being multiplied by $\prod\limits_{i=1}^{s}(t_i - 1)$, it becomes a power series, i.e. an element of $\mathbb{Z}[[t_1, \ldots, t_s]]$.

If there is a linear action (a representation) of a group G on the space A and the filtration $\{J(\underline{v})\}$ ($\underline{v} \in \mathbb{Z}^s$) is invariant with respect to this representation (i.e. $g^* J(\underline{v}) = J(\underline{v})$ for $g \in G$), one can define the equivariant Poincaré series of the filtration $\{J(\underline{v})\}$ as an element of $R(G)[[t_1, \ldots, t_s]]$. For that $d(v)$ in (1) should be substitute by $[J(\underline{v})/J(\underline{v} + \underline{1})]$ considered as a G-module (i.e. an element of the ring $R(G)$). Filtrations on the ring $\mathcal{O}_{V,0}$ considered here are, generally speaking, not G-invariant: the action of the group permutes valuations. This difficulty can be (partially) avoided by considering filtrations on the spaces of equivariant functions. This way an equivariant Poincaré series can be defined as an element of $R_1(G)[[t_1, \ldots, t_s]]$.

In the case when A is the ring $\mathcal{O}_{V,0}$ of germs of functions on $(V, 0)$ or a subspace of it, one can define a notion of integration with respect to the Euler characteristic over the projectivization $\mathbb{P}A$ of the space A (see, e.g., [2], [3]; there the notion is defined for $A = \mathcal{O}_{V,0}$, however there is no essential difference with the general case $A \subset \mathcal{O}_{V,0}$: the role of the jet spaces $J^k_{V,0} = \mathcal{O}_{V,0}/\mathfrak{m}^{k+1}$ ($\mathfrak{m}$ is the maximal ideal in the ring $\mathcal{O}_{V,0}$) is played by the spaces $JA^k_{V,0} = A/\mathfrak{m}^{k+1} \cap A$). This notion generalizes the usual notion of integration with respect to the Euler characteristic ([10]) and was inspired by the notion of motivic integration (see, e.g., [7]).

Definition: The (multi-index) filtration $\{J(\underline{v})\}$ on a subspace $A \subset \mathcal{O}_{V,0}$ is *finitely determined* if for each $\underline{v} \in \mathbb{Z}^s$ there exists $N \geq 0$ such that $\mathfrak{m}^N \cap A \subset J(\underline{v})$.

If the filtration $\{J(\underline{v})\}$ is finitely determined, one can show that

$$P(\underline{t}) = \int\limits_{\mathbb{P}A} \underline{t}^{\underline{v}(f)} d\chi \, ;$$

see, e.g., [2], [3].

Here we compute the Poincaré series for some multi-index filtrations (divisorial ones and those defined by invariant curves) on the space of equivariant function-germs on the plane $(\mathbb{C}^2, 0)$ with a finite group action. For the trivial (1-dimensional) representation of the group, this gives formulae for divisorial filtrations on the quotient surface singularity $(\mathbb{C}^2/G, 0)$ and for the filtrations defined by curves on them somewhat different from those in [3], [4].

1 Divisorial filtrations on the spaces of equivariant functions

Let $\pi : (\widetilde{V}, D) \to (V, 0)$ be a resolution of the germ $(V, 0)$. Here $\widetilde{V}$ is a smooth manifold, π is a proper analytic map which is an isomorphism outside of the exceptional divisor $D = \pi^{-1}(0)$, D is a normal crossing divisor. Let E_σ be irreducible components of the divisor D: $D = \bigcup_\sigma E_\sigma$. For a germ $f \in \mathcal{O}_{V,0}$, let $v_\sigma(f)$ be the multiplicity of the lifting $\widetilde{f} = f \circ \pi$ of the germ f to the space $\widetilde{V}$ of the resolution along the component E_σ. The function $v_\sigma : \mathcal{O}_{V,0} \setminus \{0\} \to \mathbb{Z}_{\geq 0} \cup \{+\infty\}$ defines a valuation (a *divisorial* one) on the field of quotients of the ring $\mathcal{O}_{V,0}$.

Suppose that the resolution $\pi : (\widetilde{V}, D) \to (V, 0)$ is equivariant with respect to an action of a finite group G on $(V, 0)$, i.e., that the action of G on $(V, 0)$ can be lifted to an action on $(\widetilde{V}, D)$. In particular the group G acts on the set $\{E_\sigma\}$ of irreducible components of the exceptional divisor D by permutations. If the components E_{σ_1} and E_{σ_2} are in the same orbit of the action and a function $f \in \mathcal{O}_{V,0}$ is equivariant (with respect to a 1-dimensional representation $\alpha : G \to \mathbb{C}^*$), then $v_{\sigma_1}(f) = v_{\sigma_2}(f)$.

Let us fix s components $E_1, \ldots, E_s$ of the exceptional divisor D. The corresponding valuations $v_1, \ldots, v_s$ define a multi-index filtration on the ring $\mathcal{O}_{V,0}$ of germs of functions on $(V, 0)$ and also a filtration on the space $\mathcal{O}^\alpha_{V,0}$ of functions equivariant with respect to a 1-dimensional representation α. Let $J^\alpha(\underline{v}) := \{f \in \mathcal{O}^\alpha_{V,0} : v_i(f) \geq v_i, i = 1, \ldots, s\}$. It is easy to see that the filtration $\{J^\alpha(\underline{v})\}$ is finitely determined. Let $P^\alpha(\underline{t})$ ($\underline{t} = (t_1, \ldots, t_s)$) be the Poincaré series of the filtration $\{J^\alpha(\underline{v})\}$. One has

$$P^\alpha(\underline{t}) = \int\limits_{\mathbb{P}\mathcal{O}^\alpha_{V,0}} \underline{t}^{\underline{v}(f)} d\chi \, .$$

Remarks. **1.** It is somewhat natural to suppose that for $i \neq j$ the divisors E_i and E_j are in different orbits of the action of the group G, however, from formal point of view this is not important. One can even permit that $E_i = E_j$ for some i and j.

2. In fact it is sufficient to consider the case when $\{E_i\} = \{E_\sigma\}$, i.e. $E_1, \ldots, E_s$ are all the components of the exceptional divisor D. The Poincaré series corresponding to a subset of components of the exceptional divisor is obtained from the one corresponding to all the components by putting additional variables equal to 1.

Definition: The series

$$P^G(\underline{t}) = \sum_{\alpha \in G^*} \alpha \cdot P^\alpha(\underline{t}) \in R_1(G)[[t_1, \ldots, t_s]],$$

where the sum is over all non-isomorphic 1-dimensional representations α of the group G, will be called the *equivariant Poincaré series* of the divisorial filtration corresponding to the components $E_1, \ldots, E_s$ of the exceptional divisor D of the resolution $\pi : (\widetilde{V}, D) \to (V, 0)$.

The discussion above implies that the equivariant Poincaré series $P^G(\underline{t})$ can be represented as

$$\int\limits_{\coprod_\alpha \mathbb{P}\mathcal{O}^\alpha_{V,0}} \alpha(f) \cdot \underline{t}^{\underline{v}(f)} d\chi,$$

where $\alpha : \coprod_\alpha \mathbb{P}\mathcal{O}^\alpha_{V,0} \to R_1(G)$ is the (tautological) function which sends $\mathbb{P}\mathcal{O}^\alpha_{V,0}$ to $\alpha \in R_1(G)$.

The ring $R_1(G)$ can be described in the following way. Let the abelian group $G/(G,G)$ ((G,G) is the commutator of the group G) be the direct sum $\bigoplus\limits_{j=1}^{q} \mathbb{Z}_{m_j}$ of the cyclic groups $\mathbb{Z}_{m_j} = \mathbb{Z}/m_j\mathbb{Z}$, let $\widehat{\xi_j}$ be a generator of the group $\mathbb{Z}_{m_j}$, let ξ_j be an element of the group G which maps to $\widehat{\xi_j}$ under the factorization by the commutator (G,G), and let u_j be the representation of the group G defined by $u_j(\xi_j) = \exp \frac{2\pi i}{m_j}$, $u_j(\xi_k) = 1$ for $k \neq j$. Then the representations $\prod\limits_{j=1}^{q} u_j^{\ell_j}$ with $0 \le \ell_j < m_j$ form a basis of the ring $R_1(G)$ as a $\mathbb{Z}$-module and as a ring $R_1(G) \cong \mathbb{Z}[u_1, \ldots, u_q]/(u_j^{m_j} - 1;\ j = 1, \ldots, q)$ (see e.g. [9]).

2 Divisorial filtrations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$

From now on let $(V,0) = (\mathbb{C}^2, 0)$, a finite group G acts on $(\mathbb{C}^2,0)$. Without loss of generality we suppose this action to be defined by a representation $G \to \mathbf{GL}(2,\mathbb{C})$. Let $\pi : (X, D) \to (\mathbb{C}^2, 0)$ be a modification of the plane $(\mathbb{C}^2, 0)$ by a sequence of blow-ups invariant with respect to the group action, that is the action of the group G on the plane $(\mathbb{C}^2,0)$ lifts to its action on the space X of the modification. All components E_σ of the exceptional divisor D are isomorphic to the complex projective line $\mathbb{CP}^1$. Let $\overset{\bullet}{E}_\sigma$ be the "smooth part" of the component E_σ, i.e. E_σ itself minus intersection points with all other components of the exceptional divisor D, let $\overset{\bullet}{D} = \bigcup\limits_\sigma \overset{\bullet}{E}_\sigma$ be the smooth part of the exceptional divisor D, and let $\widehat{D} = \overset{\bullet}{D}/G$ be the corresponding factor space, i.e. the space of orbits of the action of the group G on $\overset{\bullet}{D}$.

Let $-k_\sigma$ be the self intersection number $E_\sigma \circ E_\sigma$ of the component E_σ on the (smooth) surface X. Let $(E_\sigma \circ E_{\sigma'})$ be the intersection matrix of the components E_σ of the exceptional divisor D: $E_\sigma \circ E_\sigma = -k_\sigma$; for $\sigma \neq \sigma'$ one has $E_\sigma \circ E_{\sigma'} = 1$ if the components E_σ and $E_{\sigma'}$ intersect (at a point) and $E_\sigma \circ E_{\sigma'} = 0$ otherwise. One has $\det(E_\sigma \circ E_{\sigma'}) = (-1)^{\#\sigma}$ where $\#\sigma$ is the number of components of the exceptional divisor D, i.e. the number of blow-ups which produce the modification (X, D). Let $M = (m_{\sigma\sigma'}) := -(E_\sigma \circ E_{\sigma'})^{-1}$. All entries of the matrix M are positive integers. The meaning of the entry $m_{\sigma\sigma'}$ is the following. Let $\widetilde{L}_\sigma$ be a germ of a smooth curve on X transversal to the component E_σ at a smooth point of it, i.e. at a point of $\overset{\bullet}{E}_\sigma$. Let the curve $L_\sigma = \pi(\widetilde{L}_\sigma) \subset (\mathbb{C}^2, 0)$ be given by an equation $h_\sigma = 0$, $h_\sigma \in \mathcal{O}_{\mathbb{C}^2,0}$. Then $m_{\sigma\sigma'} = v_{\sigma'}(h_\sigma)$ $(= v_\sigma(h_{\sigma'}))$. For $\sigma \neq \sigma'$ the entry $m_{\sigma\sigma'}$ is also equal to the intersection number $L_\sigma \circ L_{\sigma'}$. The diagonal entry $m_{\sigma\sigma}$ is equal to $L_\sigma \circ L'_\sigma$ where $L_\sigma = \pi(\widetilde{L}_\sigma)$, $L'_\sigma = \pi(\widetilde{L}'_\sigma)$ for smooth curves $\widetilde{L}_\sigma$ and $\widetilde{L}'_\sigma$ transversal to the component E_σ at different smooth points.

For a space Z, let $S^kZ = Z^k/S_k$ be the kth symmetric power of the space Z. If Z is a quasiprojective variety, S^kZ is also one. For example $S^k\mathbb{C} \cong \mathbb{C}^k$, $S^k\mathbb{CP}^1 \cong \mathbb{CP}^k$.

Let E_0 be the first (in the order of blow-ups) component of the exceptional divisor D (E_0 may very well coincide with one of the chosen components E_1, …, E_s). The component E_0 is invariant with respect to the action of the group G. For a point p of E_0, i.e. for a line in $\mathbb{C}^2$, let ℓ_p be the corresponding linear function on $\mathbb{C}^2$: the function which vanishes at p; it is well defined up to multiplication by a non-zero constant. Let $G_p = \{g \in G : gp = p\}$ be the stabilizer of the point p and let $Gp \cong G/G_p$ be the orbit of the point p. The homogeneous function $\prod\limits_{q \in Gp} \ell_q$ of degree equal to the number $|Gp| = |G|/|G_p|$ of points in the orbit of p is an equivariant one (with respect to a certain 1-dimensional representation of the group G). Let u_p be the corresponding representation.

Examples. **1.** Let the group $\mathbb{Z}_m$ act on $(\mathbb{C}^2, 0)$ by $\xi(x, y) = (\xi^k x, \xi^\ell y)$ where $\xi = \exp(2\pi i/m)$ and $\gcd(k, \ell)$ is relatively prime with m (k and ℓ are the weights of the representation of the group $\mathbb{Z}_m$). The group $\mathbb{Z}_m^*$ of 1-dimentional representation of the group $\mathbb{Z}_m$ is the cyclic group of order m generated by the representation u defined by $u(\xi) = \xi$. The action of the group $\mathbb{Z}_m$ on the projectivization $\mathbb{CP}^1 = (\mathbb{C}^2 \setminus \{0\})/\mathbb{C}^*$ of the plane $\mathbb{C}^2$ has two fixed points $p_1 = (1, 0)$ and $p_2 = (0, 1)$ and is free outside of them. The linear function which vanishes at the point p_1 is the coordinate y. Therefore $u_{p_1} = u^{-\ell}$. In the same way $u_{p_2} = u^{-k}$. For a point $p \in \mathbb{CP}^1$ different from p_1 and p_2 one has $u_p = 1$.

2. Let $G = D_2^*$ be the subgroup of $\mathbf{SL}(\mathbb{C}, 1)$ generated by the transformations $\sigma(x, y) = (ix, -iy)$ and $\tau(x, y) = (iy, ix)$, $i = \sqrt{-1}$. The group G is of order 8 and has 3 cyclic subgroups of order 4 generated by σ, τ, and $\tau\sigma$ respectively. Intersection of any two of these subgroups is the commutator $(G, G) = \{1, \sigma^2\}$ of the group G ($\sigma^2 = \tau^2 = (\tau\sigma)^2$). The group G^* of 1-dimensional representations of the group G is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$ and is generated by the representations u_1 and u_2 defined by $u_1(\sigma) = -1$, $u_1(\tau) = 1$, $u_2(\sigma) = 1$, $u_2(\tau) = -1$; $R_1(G) = \mathbb{Z}[u_1, u_2]/(u_1^2 - 1, u_2^2 - 1)$.

The isotropy group of a generic point of the action of the group G on the projectivization $\mathbb{CP}^1 = (\mathbb{C}^2 \setminus \{0\})/\mathbb{C}^*$ consists of 2 elements and is the commutator (G, G) of the group G. Each cyclic subgroup of order 4 has two fixed points which are in the same orbit of the G-action. They are $Q_1 = (1 : 0)$ and $Q_2 = (0 : 1)$ for the subgroup $\langle\sigma\rangle$, $Q_3 = (1 : 1)$ and $Q_4 = (1 : -1)$ for the subgroup $\langle\tau\rangle$, $Q_5 = (1 : i)$ and $Q_6 = (1 : -i)$ for the subgroup $\langle\tau\sigma\rangle$. The pairs (Q_1, Q_2), (Q_3, Q_4), and (Q_5, Q_6) are orbits of the action of the group G on $\mathbb{CP}^1$.

The product of the linear functions vanishing at the points Q_1 and Q_2 is xy. It is invariant with respect to the transformation σ and anti-invariant with respect to the transformation τ. Therefore $u_{Q_1} = u_{Q_2} = u_2$. In the same way $u_{Q_3} = u_{Q_4} = u_1$, $u_{Q_5} = u_{Q_6} = u_2$ (the corresponding products of linear functions are $x^2 - y^2$ and $x^2 + y^2$ respectively). The product of linear functions vanishing at the points of an orbit of the action of the group G different from those consisting of the points Q_i is a homogeneous polynomial of degree 4 and is invariant with respect to the G-action. Therefore for any $Q \neq Q_i$, $i = 1, \ldots, 6$, one has $u_Q = 1$.

Let $\{\Xi\}$ be a stratification of the space (in fact of a smooth curve) $\widehat{D}$ ($\widehat{D} = \bigcup \Xi$) such that:

1. Each stratum Ξ is connected and therefore is contained in the image of one component E_σ under the map $j : \overset{\bullet}{D} \to \widehat{D}$ of factorization.
2. Over each strutum Ξ the map j is a covering.

As above, for a point $p \in \overset{\bullet}{D}$, let G_p be its stabilizer $\{g \in G : gp = p\}$ and let Gp be the orbit of the point p. For all points from a connected component of the preimage $j^{-1}(\Xi)$ of a stratum Ξ the stabilizer G_p is one and the same and for all points of $j^{-1}(\Xi)$ the stabilizers are conjugate to each other. For $p \in \overset{\bullet}{D}$, let $\sigma = \sigma(p)$ be the number of the component E_σ which contains the point p.

Let

$$m_{\Xi,i} \;:=\; \sum_{p\in j^{-1}(\widehat{p})} m_{\sigma(p)i} \quad \text{for } \widehat{p}\in\Xi,$$

$\underline{m}_\Xi := (m_{\Xi,1}, \ldots, m_{\Xi,s})$.

For $p \in \overset{\bullet}{D}$, let p_0 be the corresponding point of the component E_0 of the exceptional divisor: if $p \notin E_0$, p_0 is the point which belongs to the closure of the corresponding component of the complement $D \setminus E_0$. Otherwise $p_0 = p$. Let

$$u_\Xi \;:=\; u_{p_0}^{m_{\sigma(p)0}(|Gp|/|Gp_0|)}\,,$$

where $p \in j^{-1}(\Xi)$. The (1-dimensional) representation u_Ξ is a monomial, say $\underline{u}^{\underline{\ell}_\Xi}$, in basic representations, $\underline{u} = (u_1, \ldots, u_q)$, $\underline{\ell}_\Xi = (\ell_1, \ldots, \ell_q)$.

Theorem 1

$$P^G(\underline{t}) \;=\; \prod_\Xi \left(1 - \underline{u}^{\underline{\ell}_\Xi}\underline{t}^{\underline{m}_\Xi}\right)^{-\chi(\Xi)}\,. \tag{2}$$

Proof. Let us fix $\underline{V} \in \mathbb{Z}^s_{\geq 0}$ and suppose that we compute the Poincaré series $P^G(\underline{t})$ up to terms of degree $\underline{V}$ in $t_1, \ldots, t_s$. Let $\mathbb{P}\mathcal{O}^\alpha_{\mathbb{C}^2,0}(\underline{V})$ be the set $\{f \in \mathbb{P}\mathcal{O}^\alpha_{\mathbb{C}^2,0} : \underline{v}(f) \leq \underline{V}\}$. We can make additional blow-ups of intersection points of components of the exceptional divisor so that, for any $f \in \mathcal{O}_{\mathbb{C}^2,0}$ with $\underline{v}(f) \leq \underline{V}$, the strict transform of the curve $\{f = 0\}$ intersects the exceptional divisor only at smooth points. We shall keep the notation (X, D) for this modification as well explaining later why new-born components do not affect the answer. The space of G-invariant effective divisors on $\overset{\bullet}{D}$, i.e. of G-invariant unordered collection of points of $\overset{\bullet}{D}$ with some multiplicities, can be in the natural way identified with the space

$$Y^G \;=\; \prod_\Xi \left(\bigcup_{k=0}^{\infty} S^k\Xi\right) = \bigcup_{\{k_\Xi\}} \left(\prod_\Xi S^{k_\Xi}\Xi\right)\,.$$

The space Y^G can be considered as a semigroup with the union of collections as the semigroup operation. Let

$$I_G : \coprod_\alpha \mathbb{P}\mathcal{O}^\alpha_{\mathbb{C}^2,0}(\underline{V}) \to Y^G$$

be the map which sends a (non-zero) function $f \in \mathcal{O}^\alpha_{\mathbb{C}^2,0}(\underline{V})$ to the intersection of the strict transform of the zero-level curve $\{f = 0\}$ with the exceptional divisor D (a G-invariant collection of the intersection points counted with the corresponding multiplicities).

Let $\underline{v} : Y^G \to \mathbb{Z}^s_{\geq 0}$ be the map (a semigroup homomorphism) which sends a point of the component $\prod\limits_{\Xi} S^{k_\Xi}\Xi$ of the space Y^G to $\sum\limits_{\Xi} k_\Xi \underline{m}_\Xi$. Let $\underline{u} : Y^G \to G^*$ be the map to the group G^* of one-dimensional representations of the group G which sends a point of the component $\prod\limits_{\Xi} S^{k_\Xi}\Xi$ of the space Y^G to $\prod\limits_{\Xi} u_\Xi^{k_\Xi}$. One can easily see that $\underline{v} \circ I_G = \underline{v}$ and $\underline{u} \circ I_G = \underline{u}$ where $\underline{v}$ and $\underline{u}$ in the right hand sides of the equations are the corresponding maps from $\coprod\limits_{\alpha} \mathbb{P}\mathcal{O}^{\alpha}_{\mathbb{C}^2,0}(\underline{V})$. The image of the map I_G is the union of all components of the space Y^G with $\underline{v} \leq \underline{V}$. For a space (a quasi-algebraic variety) Z one has

$$\sum_{k=0}^{\infty} \chi(S^k Z) t^k = (1-t)^{-\chi(Z)}\,.$$

This implies that

$$\int\limits_{Y^G} \underline{u}\,\underline{t}^{\underline{v}} d\chi = \prod_{\Xi} \left(1 - \underline{u}^{\underline{\ell}_\Xi} \underline{t}^{\underline{m}_\Xi}\right)^{-\chi(\Xi)}\,.$$

Moreover, all additional strata corresponding to components of the exceptional divisor born under additional blow-ups have the Euler characteristics equal to zero and therefore do not participate in the right hand side of the equation.

For any point $y \in Y^G$ with $\underline{v}(y) \leq \underline{V}$, its preimage $I_G^{-1}(y)$ in $\coprod\limits_{\alpha} \mathbb{P}\mathcal{O}^{\alpha}_{\mathbb{C}^2,0}(\underline{V})$ is an affine space: [2, Proposition 2]. With the Fubini formula this implies that the integrals of $\underline{u}\,\underline{t}^{\underline{v}}$ over the spaces Y^G and $\coprod\limits_{\alpha} \mathbb{P}\mathcal{O}^{\alpha}_{\mathbb{C}^2,0}(\underline{V})$ coincide up to terms of degree $\geq \underline{V}$. Therefore Equation 2 is correct up to terms of degree $\geq \underline{V}$ for any $\underline{V} \in \mathbb{Z}^s_{\geq 0}$. This implies the statement. $\square$

3 Examples

1. Let $G = \mathbb{Z}_3$ act on the plane $(\mathbb{C}^2, 0)$ by $\xi(x, y) = (\xi x, \xi^{-1} y)$, where $\xi = \exp(2\pi i/3)$ is the generator of the group $\mathbb{Z}_3$. The factor space $(\mathbb{C}^2, 0)/\mathbb{Z}_3$ is a surface singularity of type A_2. Blowing up the origin we glue-in the exceptional divisor $\mathbb{CP}^1$ with the self-intersection number (-1). The action of the group $\mathbb{Z}_3$ on the space of the modification has two fixed points corresponding to the coordinate axes in $(\mathbb{C}^2, 0)$. At one of them (corresponding to the x-axis) the representation of the group $\mathbb{Z}_3$ has weights $(1, 1)$, at the other it has weights $(-1, -1)$. After blowing up these two points one gets a modification X of the plane $\mathbb{C}^2$ whose exceptional divisor D consists of three components with the



Figure 1: Example 1.

self-intersection numbers -1, -3, and -1. The information about the space X and the $\mathbb{Z}_3$-action on it is encoded in Figure 1.

The vertices correspond to the components of the exceptional divisor. The arrows correspond to the strict transforms of the coordinate axes in $(\mathbb{C}^2, 0)$. The numbers under the vertices are the self-intersection numbers of the corresponding components of the exceptional divisor. The numbers over each edge (mod 3) are the weights of the representation of the group $\mathbb{Z}_3$ at the corresponding intersection (fixed) point; each of these numbers corresponds to the action on the component which it is close to.

The group $\mathbb{Z}_3$ acts trivially on the first and on the third components of the exceptional divisor (counted from the left). The action on the second one is non-trivial and has two fixed points. The fact that at each fixed point of the $\mathbb{Z}_3$-action on the space X one of the weights of the representation is equal to zero implies that the factor space $X/\mathbb{Z}_3$ is smooth and is a resolution of the A_2-singularity $(\mathbb{C}^2, 0)/\mathbb{Z}_3$.

Remark. In general, for an action of a group (say, of a cyclic one) on the plane $(\mathbb{C}^2, 0)$, it is not possible to get an equivariant modification X of the plane such that the factor space X/G is non-singular and therefore is a resolution of the singularity $(\mathbb{C}^2, 0)/G$. The factor space X/G may have singular points: see e.g. Example 2 below.

One has

$$-\left(E_i \circ E_j\right) = \begin{pmatrix} 2 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}.$$

As the stratification $\{\Xi\}$ of the factor space $\widehat{D} = \overset{\bullet}{D}/\mathbb{Z}_3$ one can take the one whose three strata are isomorphic to the projective line $\mathbb{CP}^1$ minus two points (and therefore have the Euler characteristics equal to zero) and two strata consists of one point each: the images of the intersection points of the strict transforms of the coordinate axes in $\mathbb{C}^2$ with the exceptional divisor D. For these two strata the representations u_Ξ are equal to u and u^{-1} respectively.

Therefore

$$P^{\mathbb{Z}_3}(t_1, t_2, t_3) = (1 - u\,\underline{t}^{(2,1,1)})^{-1}(1 - u^{-1}\underline{t}^{(1,1,2)})^{-1}\,.$$

The exceptional divisor of the resolution $(X,0)/\mathbb{Z}_3 \to (\mathbb{C}^2,0)/\mathbb{Z}_3$ is the image of the exceptional divisor D of the modification $(X,0) \to (\mathbb{C}^2,0)$ and also consists of three components. The self-intersection numbers of these components can be determined in the following way. On the first and on the third components of the exceptional divisor D the $\mathbb{Z}_3$-action is trivial. Therefore they map isomorphically on their images. The group Z_3 acts only on the normal bundles to these components. The normal bundles to their images are the third powers of those over the preimages. Therefore the self-intersection numbers of the corresponding components of $D/\mathbb{Z}_3$ are equal to (-3). On the second component of D the $\mathbb{Z}_3$-action is not trivial. This implies that the self-intersection number of its image (the factor under the $\mathbb{Z}_3$-action) is 3 times less than the self-intersection number of the preimage and is equal to (-1). If, in $(X,0)/\mathbb{Z}_3$, we blow-down the second component of the exceptional divisor (with the self-intersection number (-1)), we get the standard (minimal) resolution of the A_2-singularity. To get the Poincaré series of the divisorial filtration corresponding to it one should take the part of the series $P^{\mathbb{Z}_3}(T_1^{1/3}, 1, T_2^{1/3})$ corresponding to the trivial representation. The reason for the change of variables $T_1 = t_1^3$, $T_2 = t_2^3$ is the difference between normal bundles to the corresponding components of D and to their images described above. This way one gets

$$P_{A_2}(T_1, T_2) = \frac{(1 - \underline{T}^{(3,3)})}{(1 - \underline{T}^{(2,1)})(1 - \underline{T}^{(1,2)})(1 - \underline{T}^{(1,1)})}$$

(cf. [3]).

2. Now let $G = \mathbb{Z}_5$ act on $(\mathbb{C}^2,0)$ by $\xi(x,y) = (\xi x, \xi^{-1}y)$ where $\xi = \exp(2\pi i/5)$. The factor space $(\mathbb{C}^2,0)/\mathbb{Z}_5$ is a surface singularity of type A_4. Blowing up the origin, after that all fixed points of the modification and then all fixed points of the second one we arrive to a modification the exceptional divisor of which has 7 irreducible components. The weights of the $\mathbb{Z}_5$-action at the fixed points which are not intersection points of the components of the exceptional divisor (they are intersection points of the exceptional divisor with the strict transforms of the coordinate axes in $\mathbb{C}^2$) are $(1,1)$ and $(-1,-1)$ respectively. Blowing up these points one gets the modification described in Figure 2.

The factor space of this modification is not smooth: it has two quotient cyclic singularities of type $(1,3)$ in the terminology of [8], corresponding to



Figure 2: Example 2.

the intersection points (edges) depicted by dashed lines. Minimal resolution of such a point has the exceptional divisor consisting of two components with the self intersection numbers -2 and -3 (see [8]). One can easily see that additional blow-ups at these points do not lead to a modification the factor space of which is smooth. Just as in Example 1 one has

$$P^{\mathbb{Z}_5}(t_1,\ldots,t_9) = (1-u\,\underline{t}^{(4,3,2,3,1,2,1,1,1)})^{-1}(1-u^{-1}\underline{t}^{(1,1,1,2,1,3,2,3,4)})^{-1}\,.$$

The components of the exceptional divisor D which correspond to the components of the standard (minimal) resolution of the A_4-singularity are those with numbers 1,4,6, and 9. One can verify this resolving the two singular points of $X/\mathbb{Z}_5$ and watching the sequence of blow-downs which leads to the minimal resolution of the A_4-singularity. On all of them the $\mathbb{Z}_5$-action is trivial and therefore they map on the four components of the resolution of the A_4-sigularity isomorphically. The Poincaré series of the divisorial filtration of (the minimal resolution of) the A_4-singularity is the part of the series $P^{\mathbb{Z}_5}(T_1^{1/5},1,1,T_2^{1/5},1,T_3^{1/5},1,1,T_4^{1/5})$ corresponding to the trivial representation. This gives

$$P_{A_4}(T_1,T_2,T_3,T_4) = \frac{(1-\underline{T}^{(5,5,5,5)})}{(1-\underline{T}^{(1,1,1,1)})\,(1-\underline{T}^{(1,2,3,4)})\,(1-\underline{T}^{(4,3,2,1)})}$$

(cf [3]).

3. Let $G = \mathbb{D}_2^*$ be the subgroup of $\mathbf{SL}(\mathbb{C},1)$ generated by the transformations $\sigma(x,y) = (ix,-iy)$ and $\tau(x,y) = (iy,ix)$, $i = \sqrt{-1}$ (see Example 2 of Section 2). The factor of the plane $\mathbb{C}^2$ by the G-action has a singular point of type D_4 (see, e.g., [8]).

Let us blow-up the origin in $\mathbb{C}^2$. On the exceptional divisor (the projective line $(\mathbb{C}^2\setminus\{0\})/\mathbb{C}^*$) there are 6 points $Q_1, \ldots, Q_6$ whose stabilizers are cyclic subgroups of G of order 4 (see Section 2, Example 2). The stabilizer of any other point of the exceptional divisor consists of 2 elements and is the commutator (G,G) of the group G. Blowing-up the points Q_i, $i=1,\ldots,6$, one gets 6 new components of the exceptional divisor. On a component corresponding to a fixed point Q_i of a cyclic subgroup H of order 4 the action of the group H is not trivial. It has two fixed points one of which is the intersection point with the exceptional divisor of the initial blow-up. The other one is the intersection point with the strict transform of the line in $\mathbb{C}^2$ corresponding to the point $Q_i \in (\mathbb{C}^2 \setminus \{0\})/\mathbb{C}^*$. Blowing-up these 6 points which do not belong to the exceptional divisor of the initial blow-up, one gets 6 new components of the exceptional divisor. On the one of them corresponding to a fixed point of a cyclic subgroup of order 4 the action of this subgroup is trivial. We arrive to the modification described in Figure 3. The arrows correspond to the strict transforms of the lines in $\mathbb{C}^2$ corresponding to the points Q_i, $i = 1, \ldots, 6$. The numbers in parenthesis are the weights of the action of the corresponding cyclic group of order 4 at the corresponding point. Let us choose one component of the exceptional divisor from each G-orbit. (The multiplicities of any equivariant function on components from the same G-orbit are equal.) These 7 components are indicated on the left hand side of Figure 3.



Figure 3: Example 3.

A stratification $\{\Xi\}$ of the factor space $\widehat{D} = \overset{\bullet}{D}/G$ can be chosen in the following way.

1. The factor space Ξ_1 of the smooth part of the initial ("central") component of the exceptional divisor. It has the Euler characteristic equal to -1. The corresponding representation u_{Ξ_1} is trivial.

2. Three strata consisting of one point each: the images of the intersection points of the exceptional divisor D with the strict transforms of the lines corresponding to the points Q_i, $i = 1, \ldots, 6$. The corresponding representations u_{Ξ_i} are: u_2 for the stratum corresponding to the fixed points of the subgroup $\langle\sigma\rangle$, u_1 for that corresponding to the subgroup $\langle\tau\rangle$, $u_1 u_2$ for that corresponding to the subgroup $\langle\tau\sigma\rangle$.

3. Six strata with the Euler characteristics equal to zero. They are images of the smooth parts of 12 components of the exceptional divisor D different from the central one minus intersection points with the strict transforms of the lines corresponding to the points Q_i.

Theorem 1 gives

$$P^G(t_1, \ldots, t_7) = \frac{1-\underline{t}^{(4,4,4,4,4,4,4)}}{(1-u_2\,\underline{t}^{(2,3,4,2,2,2,2)})(1-u_1\,\underline{t}^{(2,2,2,3,4,2,2)})(1-u_1u_2\,\underline{t}^{(2,2,2,2,2,3,4)})}\,.$$

The components of the exceptional divisor D of the modification which corresponds to the components of the standard (minimal) resolution of the D_4 singularity $(\mathbb{C}^2,0)/G$ are those with numbers 1, 3, 5, and 7. Therefore $P_{D_4}(T_1,T_2,T_3,T_4)$ is the part of the series $P^G(T_1^{1/2},1,T_2^{1/4},1,T_3^{1/4},1,T_4^{1/4})$ corresponding to the trivial representation. One gets

$$P_{D_4}(\underline{T}) = \frac{(1-\underline{T}^{(2,1,1,1)})(1+\underline{T}^{(3,2,2,2)})}{(1-\underline{T}^{(2,2,1,1)})(1-\underline{T}^{(2,1,2,1)})(1-\underline{T}^{(2,1,1,2)})}$$

(cf [3]).

4 Filtrations defined by G-invariant curves

Let $(C,0)\subset(\mathbb{C}^2,0)$ be a plane curve singularity invariant with respect to the action of the group G on $(\mathbb{C}^2,0)$. Let $C=\bigcup_{i=1}^r C_i$ be the representation of the curve C as the union of its irreducible components. Let $\varphi_i : (\mathbb{C},0)\to(\mathbb{C}^2,0)$ be the parametrization (uniformization) of the component C_i of the curve C, i.e. $\operatorname{Im}\varphi_i = C_i$, φ_i is an isomorphism between $\mathbb{C}$ and C_i outside of the origin. For a germ $f\in\mathcal{O}_{\mathbb{C}^2,0}$ let $v_i=v_i(f)$ be the power of the leading term in the power series decomposition of the germ $f\circ\varphi_i : (\mathbb{C},0)\to\mathbb{C}$: $f\circ\varphi_i(\tau)=a\tau^{v_i}+$ terms of higher degree, $a\neq 0$. If $f\circ\varphi_i(\tau)\equiv 0$, $v_i(f):=\infty$. The functions (valuations) v_i define a multi-index filtration on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ of germs of functions of two variables and also on the subspace $\mathcal{O}^{\alpha}_{\mathbb{C}^2,0}$ of functions equivariant with respect to a 1-dimensional representation α of the group G. Let $P^{\alpha}_C(t_1,\ldots,t_r)$ be the Poincaré series of this filtration on the space $\mathcal{O}^{\alpha}_{\mathbb{C}^2,0}$, let

$$P^G_C(\underline{t}) := \sum_{\alpha}\alpha\cdot P^{\alpha}_C(\underline{t})\in R_1(G)[[t_1,\ldots,t_r]]$$

(the sum is over all non-isomorphic 1-dimensional representations α of the group G) be the *equivariant Poincaré series* of the filtration on the ring $\mathcal{O}_{\mathbb{C}^2,0}$.

Remark. One can easily see that if the components C_{i_1} and C_{i_2} of the curve C are in one and the same orbit of the G-action on the set of components, $v_{i_1}(f) = v_{i_2}(f)$ for any $f \in \mathcal{O}^{\alpha}_{\mathbb{C}^2,0}$. This means that in all the monomials of the equivariant Poincaré series $P^G_C(\underline{t})$ the exponents at the variables t_{i_1} and t_{i_2} coincide. Therefore without any loss of generality one can choose one component from each orbit of the G-action.

Let $\pi : (X, D) \to (\mathbb{C}^2, 0)$ be a G-invariant embedded resolution of the curve C (i.e. the total transform $\pi^{-1}(C)$ of the curve C is a normal crossing divisor on the space X of the resolution and the group G acts on the space (a surface) X as well). Let $D = \bigcup_\sigma E_\sigma$ be the representation of the exceptional divisor $D = \pi^{-1}(0)$ as the union of its irreducible components. Let $\{\Xi\}$ be a stratification of the factor space $\widehat{D} = \overset{\bullet}{D}/G$ described in section 2. Let $\overset{\circ}{E}_\sigma$ be E_σ itself minus intersection points with all other components of the total transform $\pi^{-1}(C)$ of the curve C ($\overset{\circ}{E}_\sigma \subset \overset{\bullet}{E}_\sigma$), let $\overset{\circ}{D} = \bigcup\limits_\sigma \overset{\circ}{E}_\sigma$, and let $\widehat{D}' := \overset{\circ}{D}/G$ be the space of orbits of the action of the group G on $\widehat{D}'$. Let $\{\Xi'\}$ be a stratification of the space $\widehat{D}'$ with the same properties as the stratification $\{\Xi\}$ of the space $\widehat{D}$ in Section 2.

One has $\overset{\circ}{D} \subset \overset{\bullet}{D}$, $\widehat{D}' \subset \widehat{D}$ and one can suppose that each stratum of the stratification $\{\Xi'\}$ is a part of a stratum of the stratification $\{\Xi\}$. For a stratum Ξ', let $\underline{\ell}_{\Xi'}$ be equal to $\underline{\ell}_\Xi$ for the corresponding stratum Ξ. For $i = 1, \ldots, r$, let $E_{\sigma(i)}$ be the component of the exceptional divisor D of the resolution which intersects the component C_i of the curve C. Let $m_{\Xi',i} := m_{\Xi,\sigma(i)}$, $\underline{m}_{\Xi'} := (m_{\Xi',1}, \ldots, m_{\Xi',r})$.

Theorem 2

$$P^G_C(\underline{t}) \;=\; \prod_{\Xi'} \left(1 - \underline{u}^{\underline{\ell}_{\Xi'}} \underline{t}^{\underline{m}_{\Xi'}}\right)^{-\chi(\Xi')} .$$

The **proof** is essentially the same as the one of Theorem 1.

References

[1] Campillo A., Delgado F., Gusein-Zade S.M. The Alexander polynomial of a plane curve singularity via the ring of functions on it. Duke Math. J., v.117, no.1, 125–156 (2003).

[2] Campillo A., Delgado F., Gusein-Zade S.M. The Alexander polynomial of a plane curve singularity and integrals with respect to the Euler characteristic. Internat. J. Math., v.14, no.1, 47–54 (2003).

[3] Campillo A., Delgado F., Gusein-Zade S.M. Poincaré series of a rational surface singularity. Inventiones Math., v.155, no.1, 41–53 (2004).

[4] Campillo A., Delgado F., Gusein-Zade S.M. Poincaré series of curves on rational surface singularities. Commentarii Math. Helvetici, v.80, no.1, 95–102 (2005).

[5] Campillo A., Delgado F., Kiyek K. Gorenstein property and symmetry for one-dimensional local Cohen–Macaulay rings. Manuscripta Mathematica, v.83, no.3–4, 405–423 (1994).

[6] Delgado F., Gusein-Zade S.M. Poincaré series for several plane divisorial valuations. Proc. Edinb. Math. Soc. (2), v.46, no.2, 501–509 (2003).

[7] Denef J., Loeser F. Germs of arcs on singular algebraic varieties and motivic integration. Inventiones Math., v.135, no.1, 201–232 (1999).

[8] Pinkham, H. Singularités de Klein – I, II. In: Séminaire sur les singularités des surfaces. Lecture Notes in Math. 777, pp.1–20. Springer, Berlin–Heidelberg–New York, 1980.

[9] Serre, J.P. Représentations linéaires des groupes finis. Hermann, Paris, 1978.

[10] Viro O.Y. Some integral calculus based on Euler characteristic. Topology and Geometry – Rohlin seminar. Lecture Notes in Math. v.1346, Springer, Berlin–Heidelberg–New York, 1988, pp.127–138.

Addresses:

University of Valladolid
Dept. of Algebra, Geometry and Topology
47005 Valladolid, Spain
E-mail: campillo@cpd.uva.es, fdelgado@agt.uva.es

Moscow State University
Faculty of Mathematics and Mechanics
Moscow, 119992, Russia
E-mail: sabir@mccme.ru